

Height \_\_\_\_\_ Weight \_\_\_\_\_

## Riverbend Medical Center Regional Infusion Center (RIC) 3377 Riverbend Drive Suite 502/510 Springfield, Oregon 97477 Phone 541-222-6280 Fax 541-349-8006

## Benralizumab (FASENRA) INFUSION (v 04/18/2024)

Medicati	DIIS:
	Benralizumab 30 mg subcutaneous injection every 28 days x 3 doses then benralizumab 30 mg subcutaneous injection every 56 days
Nursing c	ommunications:
	Vital signs: Initial and prn
	Monitor patient for 30 minutes post 1st injection for hyper/ hypotension, dyspnea, nausea, itching, hives, rash, and wheezing
	Counsel patients regarding signs and symptoms of anaphylaxis with instruction to seek immediate medical care if symptoms occur
Emergen	cy Medications:
<b>■</b> diapl	DiphenhydrAMINE (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, noresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic).
mild 20 pe	- Administer 25-mg-IV-once. if-reaction-does not resolve-in 3 minutes-may repeat 25-mg-IV dose-for-a total of 50 mg-and-contact-provider to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/oints in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). act provider if given.
Avoi	Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 rd use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.
	Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.

Date: \_\_\_\_\_ Time: \_\_\_\_