

Cottage Grove Infusion 1515 Village Drive Cottage Grove, OR 97424 Phone 541-767-5447 Fax 541-767-5399 E-Fax 541-434-3164

Benralizumab (FASENRA) INFUSION (v. 05/16/2024)

Diagnosis/Indication (ICD-10): ____

Medications:

Benralizumab 30 mg subcutaneous injection every 28 days x 3 doses then benralizumab 30 mg subcutaneous injection every 56 days

Nursing communications:

- Vital signs: Initial and prn
- Monitor patient for 30 minutes post 1st injection for hyper/ hypotension, dyspnea, nausea, itching, hives, rash, and wheezing
- Counsel patients regarding signs and symptoms of anaphylaxis with instruction to seek immediate medical care if symptoms occur

Emergency Medications: (Emergency medications can be given IM if IV route unavailable)

DiphenhydrAMINE (BENADRYL) 25 to 50 mg IV as needed for mild to moderate drug reactions (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 25 mg IV once, if reaction does not resolve in 3 minutes may repeat 25 mg IV dose for a total of 50 mg and contact provider.

MethylPREDNISolone sodium succinate (Solu-MEDROL) 125 mg IV once as needed for shortness of breath, continued symptoms of mild to moderate drug reaction (flushing, dizziness, headaches, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>/= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl). Contact provider if given.

Epinephrine 0.3 mg IM once for anaphylaxis. If reaction does not resolve in 3 minutes may repeat 0.3 mg IM dose for a total of 0.6 mg. Avoid use of hand, foot, leg veins in elderly patient and those with occlusive vascular disease. Contact provider if given.

Famotidine (PEPCID) 20 mg IV once as needed for infusion/ allergic reaction.

Patient name:	Provider printed name:
DOB:	Provider signature:
Height Weight	Date: Time: