Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the  $\mathbf x$  to remove the pre-checked option.

## **ACTH STIMULATION TESTING Columbia Network Infusion Centers**

Select L	ocation: O Southwest/Vancouver C	St John/Longview	Start Date:		
Diagnos	is/Indication:				
ICD-10 (	Code(s): Auth	orization Number:			
Patient \	/itals: Height Weight				
Provi	der Communication Orders				
proces	luce delays in treatment and phone calls to y ss by selecting this option. A clinic pharmac es and procedures.				
X	I agree to utilize the PeaceHealth policies Therapeutics Committee and authorized by agreement will be issued for the duration of	y the Medical Executive C	Committee of PeaceHealth. This		
Labs		Interval			
x	ACTH Stimulation Standard Includes baseline, 30 minute, and 60 minu	Once te draws			
Supp	ortive Care	Interval			
Х	cosyntropin (CORTROSYN) injection 0.25 mg, IV, Once, Validate the baseline co	ortisol lab is collected first b	Route: IV refore Cortrosyn / ACTH is administered		
Nursing Orders		Interval			
X	Nursing Communication	Once			
	This is intended to test the patient's adrena draws are crucial to the interpretation of th methylprednisolone, or hydrocortisone in the ACTH Stim test.	e study. Please note if patie	ent has received prednisone,		
X	Nursing Communication	Once			
	Draw lab at baseline, 30 minutes, and 60 r	minutes - Cortisol Lab			
Vascu	ular Access (single select)	Interval			
0	IV	Every Visit			
	<ul> <li>Insert peripheral IV</li> <li>sodium chloride (NS) flush 10 mL</li> <li>10 mL As Needed for Line Care</li> </ul>	ŕ			
Ο	Central line (non-PICC)  - Access vascular device and confirm pa - Initiate Central line (non-PICC) mainter - sodium chloride (NS) flush 10 mL				
	10 mL As Needed for Line Care prior to medication administration PRN - sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care post medication administration PRN				
-	Provider Signature	Date	Time		
Provider	's Printed Name:				
	Place Patient Label Here	Page 1 of 2	EHR5038-DT (08/29/2024)		

- sodium chloride (NS) flush 20 mL

20 mL As Needed for Line Care post lab draw PRN

- sodium chloride (NS) flush 10 mL

10 mL for Line Care at discharge and de-access every visit

- heparin 100 units/mL 5 mL

5 mL for de-access every visit

- alteplase (CATHFLO) injection 2 mg

Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

O PICC line Every Visit

- Access vascular device and confirm patency
- Initiate PICC maintenance protocol
- Nursing Communication

Change PICC line dressing weekly and PRN

- sodium chloride (NS) flush 10 mL

10 mL As Needed for Line Care prior to medication administration PRN

sodium chloride (NS) flush 10 mL

10 mL As Needed for Line Care post medication administration PRN

- sodium chloride (NS) flush 20 mL

20 mL As Needed for Line Care post lab draw PRN

- alteplase (CATHFLO) injection 2 mg

Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

Provider Signature	Date	Time	
Provider's Printed Name:			