



Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the x to remove the pre-checked option.

DAPTOMYCIN INFUSION [11500420] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.

Labs	Interval
<input type="radio"/> Complete Blood Count W/ Automated Differential	Frequency _____
<input type="radio"/> Comprehensive Metabolic Panel	Frequency _____
<input type="radio"/> Creatine Kinase, Total	Frequency _____
<input type="radio"/> Sedimentation Rate, Westergren	Frequency _____
<input type="radio"/> C-Reactive Protein	Frequency _____

Supportive Care	Interval
<input checked="" type="checkbox"/> DAPTOMYCIN Dose _____ mg / Dose Basis _____ mg/kg	Route: IV Frequency _____
<input checked="" type="checkbox"/> DURATION / END DATE: _____	

Nursing Orders	Interval
<input type="radio"/> Nursing Communication <i>RN to notify provider if CPK > 1000 units/L with unexplained signs and symptoms of myopathy and/or if CPK > 2000 units/L.</i>	Every 7 Days
<input checked="" type="checkbox"/> Nursing Communication <i>At the end of treatment, contact MD/provider to address removal of PICC line.</i>	Every Visit
<input checked="" type="checkbox"/> Nursing Communication <i>Discontinue therapy plan when treatment complete.</i>	Every Visit

Vascular Access (single select)	Interval
<input type="radio"/> IV - Insert peripheral IV - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	Every Visit

Provider Signature _____ Date _____ Time _____

Provider's Printed Name: _____

Place Patient Label Here

- Central line (non-PICC) Every Visit
 - Access vascular device and confirm patency
 - Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
5 mL for de-access every visit
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

- PICC line Every Visit
 - Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications	Interval	
○ sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	PRN	Route: IV
○ sodium chloride 0.9 % bolus 500 mL <i>500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)</i>	PRN	Route: IV

Provider Signature Date Time

Provider's Printed Name: