



Change order details by crossing out unwanted information and writing in desired details/instructions.
Place a line through the x to remove the pre-checked option.

INFLIXIMAB OR BIOSIMILARS - INITIATION DOSES (0 , 2 , & 6 WEEKS) [11500955] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.**

Pre-Screening Labs

- Provider Communication
Provider to send PPD/Tuberculosis screening and Hep B screening lab results with completed orders.

Labs

Interval

- Complete Blood Count W/ Automated Differential Frequency _____
- Comprehensive Metabolic Panel Frequency _____

Pre-Medications

Interval

- | | | |
|--|-------------|-------------|
| <input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg | Every Visit | Route: Oral |
| <input type="checkbox"/> loratadine (CLARITIN) tablet 10 mg | Every Visit | Route: Oral |
| <input type="checkbox"/> diphenhydrAMINE (BENADRYL) capsule 25 mg | Every Visit | Route: Oral |
| <input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 40 mg | Every Visit | Route: IV |

Supportive Care

Interval

- | | |
|---|-----------------|
| <input type="checkbox"/> INFLIXIMAB IVPB (REMICADE) | Route: IV |
| <i>Starting when released, For 1 dose, DOSING WILL BE ROUNDED TO NEAREST VIAL SIZE. Use an in-line, sterile, non-pyrogenic, low protein-binding filter with 1.2 micron pore size or less. Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i> | |
| Dose _____ | Frequency _____ |
| <input type="checkbox"/> INFLIXIMAB-DYYB IVPB (INFLECTRA) | Route: IV |
| <i>Starting when released, For 1 dose, DOSING WILL BE ROUNDED TO NEAREST VIAL SIZE. Use an in-line, sterile, non-pyrogenic, low protein-binding filter with 1.2 micron pore size or less. Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i> | |
| Dose _____ | Frequency _____ |
| <input type="checkbox"/> INFLIXIMAB-ABDA IVPB (RENFLEXIS) | Route: IV |
| <i>Starting when released, For 1 dose, DOSING WILL BE ROUNDED TO NEAREST VIAL SIZE. Use an in-line, sterile, non-pyrogenic, low protein-binding filter with 1.2 micron pore size or less. Infuse 10ml/hr x15min; then 20ml/hr x15min; then 40ml/hr x15min; then 80ml/hr x15min; then 150ml/hr x15min; then max 250ml/hr until done.</i> | |
| Dose _____ | Frequency _____ |

Provider Signature

Date

Time

Provider's Printed Name: _____

Place Patient Label Here

Page 1 of 3

EHR2028-DT (07/10/2024)

**INFLIXIMAB OR BIOSIMILARS - INITIATION DOSES
(0 , 2 , & 6 WEEKS)**

Nursing Orders		Interval
<input checked="" type="checkbox"/>	Nursing Communication <i>Assess patients' vital signs prior to the infliximab infusion, before each rate increase, and 30 minutes following the infusion. Observe patient 30 minutes after completion of infusion.,</i>	Every Visit

Vascular Access (single select)		Interval
<input type="checkbox"/>	IV - Insert peripheral IV - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	Every Visit
<input type="checkbox"/>	Central line (non-PICC) - Access vascular device and confirm patency - Initiate Central line (non-PICC) maintenance protocol - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care prior to medication administration PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care post medication administration PRN</i> - sodium chloride (NS) flush 20 mL <i>20 mL As Needed for Line Care post lab draw PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL for Line Care at discharge and de-access every visit</i> - heparin 100 units/mL 5 mL <i>5 mL for de-access every visit</i> - alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i>	Every Visit
<input type="checkbox"/>	PICC line - Access vascular device and confirm patency - Initiate PICC maintenance protocol - Nursing Communication <i>Change PICC line dressing weekly and PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care prior to medication administration PRN</i> - sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care post medication administration PRN</i> - sodium chloride (NS) flush 20 mL <i>20 mL As Needed for Line Care post lab draw PRN</i> - alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i>	Every Visit

PRN Medications		Interval	
<input type="checkbox"/>	sodium chloride (NS) flush 10 mL <i>10 mL As Needed for Line Care</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % bolus 500 mL <i>500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)</i>	PRN	Route: IV

Provider Signature _____ Date _____ Time _____

Provider's Printed Name: _____

Place Patient Label Here	Page 2 of 3 EHR2028-DT (07/10/2024) INFLIXIMAB OR BIOSIMILARS - INITIATION DOSES (0 , 2, & 6 WEEKS)
--------------------------	--

Emergency Medications**Interval**

- | X | | PRN | Route: IV |
|----------|---|------------|-----------------------------|
| | diphenhydrAMINE (BENADRYL) injection 25-50 mg
<i>25-50 mg Once As Needed For mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction. Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg I dose for a total of 50 mg, and contact provider.</i> | | |
| | albuterol 90 mcg/actuation inhaler 2 puff
<i>2 puff Once As Needed Inhalation Wheezing, Shortness of Breath, wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available.</i> | PRN | Route: Inhalation |
| | methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 125 mg
<i>125 mg Once As Needed For shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl), and contact provider.</i> | PRN | Route: IV |
| | EPINEPHrine (ADRENALIN) injection for Allergic Reaction 0.5 mg
<i>0.5 mg Once As Needed For severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (>= 40 points in SBP), shortness of breath with wheezing and O2Sat <90%), and contact provider.</i> | PRN | Route: Intramuscular |

 Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 3 of 3

EHR2028-DT (07/10/2024)

**INFLIXIMAB OR BIOSIMILARS - INITIATION DOSES
(0 , 2, & 6 WEEKS)**