



Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the **x** to remove the pre-checked option.

IRON - FERRIC CARBOXYMALTOSE (INJECTAFER) IVPB [11500556] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.**

Labs	Interval
------	----------

- | | | |
|---|-------------------------|------|
| <input checked="" type="checkbox"/> | Physician Communication | Once |
| <i>Provider to send most recent Ferritin and/or Iron Deficiency Panel Lab results with completed order.</i> | | |

Supportive Care	Interval
-----------------	----------

- | | | |
|--|---|--------------------|
| <input checked="" type="checkbox"/> | FERRIC CARBOXYMALTOSE IVPB 750 mg IV every 7 days x 2 doses | Route: IV |
| <i>Monitor patient for signs and symptoms of hypersensitivity DURING INFUSION AND FOR AT LEAST 30 MINUTES AFTER INFUSION. Hypersensitivity symptoms may include: anaphylaxis, flushing, dyspnea, tachycardia, and increased blood pressure. For patients weighing less than 50 kg, the dose will be updated to 15 mg/kg.</i> | | |
| <input checked="" type="checkbox"/> | SODIUM CHLORIDE 0.9 % IV BOLUS | PRN Route: IV |
| <i>IV site discomfort May use 250 ml NS at 100ml/hr. Run concurrent with iron infusions as needed.</i> | | |

Nursing Orders	Interval
----------------	----------

- | | | |
|---|-----------------------|--------------|
| <input checked="" type="checkbox"/> | Nursing Communication | Every 7 days |
| <i>Monitor patient for signs and symptoms of hypersensitivity DURING INFUSION AND FOR AT LEAST 30 MINUTES AFTER INFUSION. Hypersensitivity symptoms may include: anaphylaxis, flushing, dyspnea, tachycardia, and increased blood pressure.</i> | | |
| <input checked="" type="checkbox"/> | Nursing Communication | Every visit |
| <i>Discontinue therapy plan when treatment complete.</i> | | |

Vascular Access (single select)	Interval
---------------------------------	----------

- | | | |
|---|-------------------------|-------------|
| <input type="radio"/> | IV | Every Visit |
| <ul style="list-style-type: none"> - Insert peripheral IV - sodium chloride (NS) flush 10 mL <li style="padding-left: 40px;">10 mL As Needed for Line Care | | |
| <input type="radio"/> | Central line (non-PICC) | Every Visit |
| <ul style="list-style-type: none"> - Access vascular device and confirm patency | | |

Provider Signature	Date	Time
--------------------	------	------

Provider's Printed Name: _____

Place Patient Label Here

- Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
5 mL for de-access every visit
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- O** PICC line Every Visit
- Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications	Interval	Route
-----------------	----------	-------

- | | | |
|---|-----|-----------|
| <input type="radio"/> sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care | PRN | Route: IV |
| <input type="radio"/> sodium chloride 0.9 % bolus 500 mL
500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr) | PRN | Route: IV |

Emergency Medications	Interval	Route
-----------------------	----------	-------

- | | | |
|---|-----|-----------|
| <input checked="" type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25-50 mg
25-50 mg Once As Needed For mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritic). Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction. Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25 mg I dose for a total of 50 mg, and contact provider. | PRN | Route: IV |
|---|-----|-----------|

Provider Signature	Date	Time
--------------------	------	------

Provider's Printed Name:

Place Patient Label Here

Page 2 of 3

EHR5002-DT (05/24/2024)

IRON - FERRIC CARBOXYMALTOSE (INJECTAFER) IVPB

- X albuterol 90 mcg/actuation inhaler 2 puff PRN Route: Inhalation
2 puff Once As Needed Inhalation Wheezing, Shortness of Breath, wheezing, shortness of breath associated with infusion reaction and notify provider. Administer with a spacer if available.

- X methylPREDNISolone sodium succinate (Solu-MEDROL) Injection 125 mg PRN Route: IV
125 mg Once As Needed For shortness of breath for continued symptoms of mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (≥ 20 points in SBP), nausea, urticaria, chills, pruritic) that worsen or persist 5 minutes after administration of diphenhydramine (Benadryl), and contact provider.

- X EPINEPHrine (ADRENALIN) injection for Allergic Reaction 0.5 mg PRN Route: Intramuscular
0.5 mg Once As Needed For severe drug reaction (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort plus blood pressure changes (≥ 40 points in SBP), shortness of breath with wheezing and O2Sat $<90\%$), and contact provider.

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here