



Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the x to remove the pre-checked option.

LINE CARE PLAN [11500983] Columbia Network Infusion Centers

Select Location: Southwest/Vancouver St John/Longview Start Date: _____

Diagnosis/Indication: _____

ICD-10 Code(s): _____ Authorization Number: _____

Patient Vitals: Height _____ Weight _____

Provider Communication Orders

To reduce delays in treatment and phone calls to your office, you may participate in the PeaceHealth formulary process by selecting this option. A clinic pharmacist will adjust orders according to PeaceHealth site approved policies and procedures.

- I agree to utilize the PeaceHealth policies & procedures that have been reviewed by the Pharmacy & Therapeutics Committee and authorized by the Medical Executive Committee of PeaceHealth. This agreement will be issued for the duration of active orders contained within this treatment plan.

Vascular Access (single select) Interval

- Central line (non-PICC) Every Visit
 - Access vascular device and confirm patency
 - Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
5 mL for de-access every visit
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded
- PICC line Every Visit
 - Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN

Provider Signature _____ Date _____ Time _____

Provider's Printed Name: _____

Place Patient Label Here

- alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications	Interval	
○ sodium chloride (NS) flush 10 mL 10 mL As Needed for Line Care	PRN	Route: IV
○ sodium chloride 0.9 % bolus 500 mL 500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)	PRN	Route: IV

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 2 of 2

EHR5017-DT (05/24/2024)

LINE CARE PLAN