

- Central line (non-PICC) Every Visit
 - Access vascular device and confirm patency
 - Initiate Central line (non-PICC) maintenance protocol
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - sodium chloride (NS) flush 10 mL
10 mL for Line Care at discharge and de-access every visit
 - heparin 100 units/mL 5 mL
5 mL for de-access every visit
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

- PICC line Every Visit
 - Access vascular device and confirm patency
 - Initiate PICC maintenance protocol
 - Nursing Communication
Change PICC line dressing weekly and PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care prior to medication administration PRN
 - sodium chloride (NS) flush 10 mL
10 mL As Needed for Line Care post medication administration PRN
 - sodium chloride (NS) flush 20 mL
20 mL As Needed for Line Care post lab draw PRN
 - alteplase (CATHFLO) injection 2 mg
Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded

PRN Medications

Interval

- | | | |
|--|-----|-----------|
| ○ sodium chloride (NS) flush 10 mL
<i>10 mL As Needed for Line Care</i> | PRN | Route: IV |
| ○ sodium chloride 0.9 % bolus 500 mL
<i>500 mL Once for Line Care, Starting when released, For 1 dose, Run at TKO (25ml/hr)</i> | PRN | Route: IV |

Provider Signature

Date

Time

Provider's Printed Name:

Place Patient Label Here

Page 2 of 2

EHR5048-DT (08/29/2024)

PENICILLIN INFUSION