

<input type="checkbox"/>	Access & Use Central Line/CVAD <i>Starting when released Access vascular device and confirm patency.</i>	PRN	
<input type="checkbox"/>	Access & Use Central Line/CVAD <i>Starting when released Access vascular device and confirm patency.</i>	PRN	
<input type="checkbox"/>	sodium chloride (NS) flush 10 mL <i>Line Care, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	Initiate Central Line (Non-PICC) Maintenance Protocol <i>Starting when released</i>	PRN	
<input type="checkbox"/>	Initiate PICC Maintenance Protocol <i>Starting when released</i>	PRN	
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care prior to medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	Nursing Communication <i>Starting when released Change PICC line dressing weekly and as needed.</i>	PRN	
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care post medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care prior to medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 20 mL <i>Other, Line Care post lab draw, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care post medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care at discharge and de-access, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 20 mL <i>Other, Line Care post lab draw, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	heparin, porcine (PF) 100 unit/mL flush 500 Units <i>Line Care, for de-access, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	alteplase (CATHFLO) injection 2 mg <i>Other, Line Care, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded.</i>	PRN	Route: Intra-Catheter
<input type="checkbox"/>	alteplase (CATHFLO) injection 2 mg <i>Other, Line care, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i>	PRN	Route: Intra-Catheter
Pre-Medications		Interval	
<input type="checkbox"/>	acetaminophen (TYLENOL) tablet 650 mg <i>Once Starting when released, For 1 dose</i>	Every visit	Route: Oral

Provider Signature

EHR User ID

Date

Time

Initials

Place Patient Label Here

