

Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the \boxtimes to remove the pre-checked option.

METHYLPREDNISOLONE INFUSION (3-DAY) [11500398] Therapy Plan To Be Used In Infusion Center

Infusion Center Location:		Start Date:				
Diagno	sis/Indication:					
Authori	zation Number:					
Patient Name				Weight		
Supp	ortive Care METHYLPREDNISOLONE SODIUM SUCCINA		Route:			
	Starting when released, For 1 dose, Day 1		Noute.			
	Dose Frequency _	013				
Provi	der Communication Orders	Interval				
	Provider Communication	Once				
	Starting when released Select most approp central line access set of orders, then de-se		/) for IV access / cen	tral line - if selecting a		
Nursi	ng Orders	Interval				
	Insert peripheral IV	PRN				
	Starting when released					
	Access & Use Central Line/CVAD	PRN				
	Starting when released Access vascular device and confirm patency.					
	Access & Use Central Line/CVAD	PRN				
	Starting when released Access vascular device and confirm patency.					
	sodium chloride (NS) flush 10 mL	PRN	Route: IV			
	Line Care, Starting when released					
	Initiate Central Line (Non-PICC) Maintenance I	Protocol PRN				
	Starting when released					
	Initiate PICC Maintenance Protocol	PRN				
	Starting when released					
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care prior to medication administration, Starting when released					
	Nursing Communication	PRN				
	Starting when released Change PICC line of					
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care post medication administration, Starting when released					
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care prior to medication admini	-				
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV			
	Other, Line Care post lab draw, Starting wh	en released				
	Provider Signature	EHR User ID Date)	Time		
Initials		L				
		Page 1 of 2		4-DT (01/05/2024)		
Place Patient Label Here		METHYLPRE	METHYLPREDNISOLONE INFUSION (3-DAY)			
		Prog & Orders				

	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV				
	Other, Line Care post medication administration, Starting when released						
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV				
	Other, Line Care at discharge and de-access, Starting when released						
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV				
	Other, Line Care post lab draw, Starting when released						
	heparin, porcine (PF) 100 unit/mL flush 500 Units	PRN	Route: IV				
	Line Care, for de-access, Starting when released						
	alteplase (CATHFLO) injection 2 mg	PRN	Route: Intra-Catheter				
	Other, Line Care, Starting when released, For 2 do	<i>,</i> 0					
	Other, Line Care, Starting when released, For 2 do water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m	ed to allow large	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho	ed to allow large	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m	bed to allow large build occur within 3 inutes to 2 hours PRN se, For clearing co allow large bubblo occur within 3 min	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate es to dissipate. Mix by gently swirling until jutes); do not shake. Final concentration:				
	 water for injection to vial; let the vial stand undisturbuttly until completely dissolved (complete dissolution shock concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should be completed) 	bed to allow large build occur within 3 inutes to 2 hours PRN se, For clearing co allow large bubblo occur within 3 min	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate es to dissipate. Mix by gently swirling until jutes); do not shake. Final concentration:				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should of 1 mg/mL. Retain in catheter for 30 minutes to 2 hou	bed to allow large build occur within 3 inutes to 2 hours PRN se, For clearing co allow large bubbl occur within 3 min irs, instill a 2nd do	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate es to dissipate. Mix by gently swirling until jutes); do not shake. Final concentration:				
PRN	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should of 1 mg/mL. Retain in catheter for 30 minutes to 2 hou Medications	ped to allow large buld occur within 3 inutes to 2 hours, PRN se, For clearing co allow large bubbl occur within 3 min rrs, instill a 2nd do Interval PRN	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate es to dissipate. Mix by gently swirling until utes); do not shake. Final concentration: ose if occluded				
PRN	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should of 1 mg/mL. Retain in catheter for 30 minutes to 2 hou Medications sodium chloride (NS) flush 10 mL	ped to allow large buld occur within 3 inutes to 2 hours, PRN se, For clearing co allow large bubbl occur within 3 min rrs, instill a 2nd do Interval PRN	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate es to dissipate. Mix by gently swirling until utes); do not shake. Final concentration: ose if occluded				

	Provider Signature	EHR User ID	Date	Time	
Initials					
		Page 2 of 2		EHR0234-DT (01/05/2024)	
		METHYLPREDNISOLONE INFUSION (3-DAY)			
F	Place Patient Label Here	Prog & Orders			