

Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the \boxtimes to remove the pre-checked option.

METHYLPREDNISOLONE INFUSION (5-DAY) [11500295] Therapy Plan To Be Used In Infusion Center

Infusion Center Location:		Start Date:				
Diagno	sis/Indication:					
Author	ization Number:					
Patient Name				Weight		
			0			
Supp	ortive Care METHYLPREDNISOLONE SODIUM SUCCINA		Route:			
	Starting when released, For 1 dose, Day 1 do					
	Dose Frequency					
Provi	der Communication Orders	Interval				
	Provider Communication	Once				
	Starting when released Select most approp central line access set of orders, then de-se		y) for IV access / cen	tral line - if selecting a		
Nursi	ing Orders	Interval				
	Insert peripheral IV	PRN				
	Starting when released					
	Access & Use Central Line/CVAD	PRN				
	Starting when released Access vascular device and confirm patency.					
	Access & Use Central Line/CVAD	PRN				
	Starting when released Access vascular device and confirm patency.					
	sodium chloride (NS) flush 10 mL	PRN	Route: IV			
	Line Care, Starting when released					
	Initiate Central Line (Non-PICC) Maintenance I	Protocol PRN				
	Starting when released					
	Initiate PICC Maintenance Protocol	PRN				
	Starting when released					
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care prior to medication admini	istration, Starting when re	eleased			
	Nursing Communication	PRN				
	Starting when released Change PICC line of	dressing weekly and as r	needed.			
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care post medication administra	ation, Starting when rele	ased			
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care prior to medication admini	istration, Starting when re	eleased			
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV			
	Other, Line Care post lab draw, Starting when released					
	Provider Signature	EHR User ID Date	e	Time		
Initials						
		Page 1 of 2	EHR023	5-DT (01/05/2024)		
Place Patient Label Here		-		· · · · ·		
l		Prog & Orders				

	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV				
	Other, Line Care post medication administration, Starting when released						
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV				
	Other, Line Care at discharge and de-access, Starting when released						
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV				
	Other, Line Care post lab draw, Starting when released						
	heparin, porcine (PF) 100 unit/mL flush 500 Units	PRN	Route: IV				
	Line Care, for de-access, Starting when released						
	alteplase (CATHFLO) injection 2 mg	PRN	Route: Intra-Catheter				
	Other, Line Care, Starting when released, For 2 do	<i>,</i> 0					
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m	bed to allow large buld occur within inutes to 2 hours	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded.				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho	ed to allow large	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m	bed to allow large build occur within s inutes to 2 hours PRN se, For clearing c allow large bubbl occur within 3 min	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wat les to dissipate. Mix by gently swirling until nutes); do not shake. Final concentration: 1				
	 water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to a completely dissolved (complete dissolution should completely dissolved (complete dissolution should complete dissolution should complete dissolution should completely dissolved (complete dissolution should complete dissolution should complete	bed to allow large build occur within s inutes to 2 hours PRN se, For clearing c allow large bubbl occur within 3 min	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wat les to dissipate. Mix by gently swirling until nutes); do not shake. Final concentration: 1				
	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should c mg/mL. Retain in catheter for 30 minutes to 2 hours	bed to allow large buld occur within S inutes to 2 hours PRN se, For clearing c allow large bubbl occur within 3 min s, instill a 2nd dos	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wat les to dissipate. Mix by gently swirling until nutes); do not shake. Final concentration: 1				
PRN	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to completely dissolved (complete dissolution should of mg/mL. Retain in catheter for 30 minutes to 2 hours Medications	bed to allow large buld occur within s inutes to 2 hours PRN se, For clearing c allow large bubbl occur within 3 min s, instill a 2nd dos Interval PRN	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate les to dissipate. Mix by gently swirling until nutes); do not shake. Final concentration: 1 e if occluded				
PRN	water for injection to vial; let the vial stand undisturb until completely dissolved (complete dissolution sho concentration: 1 mg/mL. Retain in catheter for 30 m alteplase (CATHFLO) injection 2 mg Other, Line care, Starting when released, For 2 dos for injection to vial; let the vial stand undisturbed to a completely dissolved (complete dissolution should c mg/mL. Retain in catheter for 30 minutes to 2 hours Medications sodium chloride (NS) flush 10 mL	bed to allow large buld occur within s inutes to 2 hours PRN se, For clearing c allow large bubbl occur within 3 min s, instill a 2nd dos Interval PRN	bubbles to dissipate. Mix by gently swirling 3 minutes); do not shake. Final , instill a 2nd dose if occluded. Route: Intra-Catheter entral line catheter. Add 2.2 mL sterile wate les to dissipate. Mix by gently swirling until nutes); do not shake. Final concentration: 1 e if occluded				

	Provider Signature	EHR User ID	Date	Time	
Initials					
		Page 2 of 2		EHR0235-DT (01/05/2024)	
		METHYLPREDNISOLONE INFUSION (5-DAY)			
F	Place Patient Label Here	Prog & Orders			