

Place Patient Label Here

Change order details by crossing out unwanted information and writing in desired details/instructions. Place a line through the  $\boxtimes$  to remove the pre-checked option.

## METHYLPREDNISOLONE INFUSION (Q2WK) [11500400] Therapy Plan To Be Used In Infusion Center

Infusion Center Location:		Start Date	e:					
Diagnosis/Indication:								
Authorization Number:								
Patient Name		_ DOB	Height Weig	ht				
Supportive Care		Interval						
	METHYLPREDNISOLONE SODIUM SUCCINATION	TE IV/IM ORDERABLE	Route:					
	Starting when released, For 1 dose Dose Frequency _							
Provider Communication Orders		Interval						
	Provider Communication	Once						
	Starting when released Select most appropriate orders (one set only) for IV access / central line - if selecting central line access set of orders, then de-select IV access.							
Nursi	ing Orders	Interval						
	Insert peripheral IV	PRN						
	Starting when released							
	Access & Use Central Line/CVAD	PRN						
	Starting when released Access vascular de	vice and confirm patenc	у.					
	Access & Use Central Line/CVAD	PRN						
	Starting when released Access vascular de	vice and confirm patenc	у.					
	sodium chloride (NS) flush 10 mL	PRN	Route: IV					
	Line Care, Starting when released							
	Initiate Central Line (Non-PICC) Maintenance F	Protocol PRN						
	Starting when released							
	Initiate PICC Maintenance Protocol	PRN						
	Starting when released							
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV					
	Other, Line Care prior to medication adminis	er, Line Care prior to medication administration, Starting when released						
	Nursing Communication	PRN						
	Starting when released Change PICC line dressing weekly and as needed.							
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV					
	Other, Line Care post medication administration, Starting when released							
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV					
	eleased							
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV					
Other, Line Care post lab draw, Starting when released								
	Provider Signature	EHR User ID Dat	e Time					
Initials								
		Page 1 of 2	EHR0668-DT (0 <sup>-</sup>	1/05/2024)				

METHYLPREDNISOLONE INFUSION (Q2WK)

Prog & Orders

	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care post medication administration, Starting when released					
	sodium chloride 0.9 % injection 10 mL	PRN	Route: IV			
	Other, Line Care at discharge and de-access, Starting	when released				
	sodium chloride 0.9 % injection 20 mL	PRN	Route: IV			
	Other, Line Care post lab draw, Starting when released	d				
	heparin, porcine (PF) 100 unit/mL flush 500 Units	PRN	Route: IV			
	Line Care, for de-access, Starting when released					
	alteplase (CATHFLO) injection 2 mg	PRN	Route: Intra-Catheter			
	Other, Line Care, Starting when released, For 2 dose, water for injection to vial; let the vial stand undisturbed until completely dissolved (complete dissolution should concentration: 1 mg/mL. Retain in catheter for 30 minut	to allow large bubl occur within 3 mir	bles to dissipate. Mix by gently swirling nutes); do not shake. Final			
	alteplase (CATHFLO) injection 2 mg	PRN	Route: Intra-Catheter			
	Other, Line care, Starting when released, For 2 dose, I for injection to vial; let the vial stand undisturbed to allo completely dissolved (complete dissolution should occumg/mL. Retain in catheter for 30 minutes to 2 hours, in:	w large bubbles to ır within 3 minutes	dissipate. Mix by gently swirling until ); do not shake. Final concentration: 1			
PRN Medications		Interval				
	sodium chloride (NS) flush 10 mL	PRN	Route: IV			
	10 mL As Needed IV Line Care, Starting when release	d				
	sodium chloride 0.9 % bolus 500 mL	PRN	Route: IV			
	500 mL Once As Needed IV Line Care, Starting when released, Run at TKO (25ml/hr)					

Provider Signature	EHR User ID	Date	Time
Initials			

Page 2 of 2

EHR0668-DT (01/05/2024)

METHYLPREDNISOLONE INFUSION (Q2WK)

Place Patient Label Here

Prog & Orders