

OP Rabies Exposure Smart Set [3040000848]

General

Nursing

<input checked="" type="checkbox"/> Weigh patient	Routine, Once, Starting Today For 1 Occurrences, OP Blood Infusion
<input checked="" type="checkbox"/> Nurse Communication - Rabies Vaccine Information	Routine, Until discontinued, Starting Today Provide patient / patient support person with CDC Rabies Vaccine Information Sheet., OP Blood Infusion

Referrals - Infusion Center

Place referral order if patient to receive medications in the infusion center (outpatient).

<input type="checkbox"/> Ambulatory referral to Infusion Therapy	Internal Referral
--	-------------------

Medications

Rabies Immune Globulin (Single Response)

Order Rabies Immune Globulin if not previously vaccinated*.

*Previously vaccinated persons are those who have received one of the recommended pre-exposure or post exposure regimens of HDCV, RA or PCEC, or those who received another vaccine and had a documented antibody titer (see ACIP Human Rabies Prevention guidelines for details.)

<input type="checkbox"/> Rabies Immune Globulin - Wound Present	
<input checked="" type="checkbox"/> rabies immune globulin (PF) (HyperRAB) injection	20 Units/kg, Intramuscular, Once, For 1 Doses, OP Blood Infusion Infiltrate as much RIG intradermally into and around wound(s) as possible. Inject remaining RIG intramuscularly in site remote from wound, preferably deltoid muscle of upper arm or lateral thigh (adults). Use anterolateral thigh in infants and small children. **Administer at an anatomical site distant from vaccine **
<input checked="" type="checkbox"/> rabies vaccine (IMOVAX/RABAVERT) injection	2.5 Units, Intramuscular, Once, For 1 Doses, OP Blood Infusion Inject vaccine at anatomical site distant from where RIG was administered. Use deltoid in adults and adolescents; anterolateral thigh in infants and small children.
<input type="checkbox"/> Rabies Immune Globulin - No Wound Present	
<input checked="" type="checkbox"/> rabies immune globulin (PF) (HyperRAB) injection	20 Units/kg, Intramuscular, Once, For 1 Doses, OP Blood Infusion Inject in gluteal sites in adults; anterolateral thigh in infants and small children, in divided doses. **Administer at an anatomical site distant from vaccine **
<input checked="" type="checkbox"/> rabies vaccine (IMOVAX/RABAVERT) injection	2.5 Units, Intramuscular, Once, For 1 Doses, OP Blood Infusion Inject vaccine at anatomical site distant from where RIG was administered. Use deltoid in adults and adolescents; anterolateral thigh in infants and small children.

Rabies Vaccine

A person who is exposed and has never been vaccinated against rabies should get 4 doses of rabies vaccine—one dose right away, and additional doses on the 3rd, 7th, and 14th days. (If previously vaccinated with HDCV, RA or PCEC or if documented antibody response, only two doses are needed: day 0 and day 3.)

(For persons with broadly defined immunosuppression, administer 5th dose of vaccine on Day 28 per ACIP Provisional Recommendations dated 6/24/09)

<input type="checkbox"/> rabies vaccine (IMOVAX/RABAVERT) injection	2.5 Units, Intramuscular, Once, For 1 Doses, OP Blood Infusion Use deltoid in adults and adolescents; anterolateral thigh in infants and small children.
---	---

Provider Signature

EHR User ID

Date

Time

Initials

Place patient label here

