

<input type="checkbox"/>	Initiate PICC Maintenance Protocol <i>Starting when released</i>	PRN	
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care prior to medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	Nursing Communication <i>Starting when released Change PICC line dressing weekly and as needed.,</i>	PRN	
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care post medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care prior to medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 20 mL <i>Other, Line Care post lab draw, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care post medication administration, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 10 mL <i>Other, Line Care at discharge and de-access, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % injection 20 mL <i>Other, Line Care post lab draw, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	heparin, porcine (PF) 100 unit/mL flush 500 Units <i>Line Care, for de-access, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded.</i>	PRN	Route: Intra-Catheter
<input type="checkbox"/>	alteplase (CATHFLO) injection 2 mg <i>Other, Starting when released, For 2 dose, For clearing central line catheter. Add 2.2 mL sterile water for injection to vial; let the vial stand undisturbed to allow large bubbles to dissipate. Mix by gently swirling until completely dissolved (complete dissolution should occur within 3 minutes); do not shake. Final concentration: 1 mg/mL. Retain in catheter for 30 minutes to 2 hours, instill a 2nd dose if occluded</i>	PRN	Route: Intra-Catheter
PRN Medications		Interval	
<input type="checkbox"/>	sodium chloride (NS) flush 10 mL <i>10 mL As Needed IV Line Care, Starting when released</i>	PRN	Route: IV
<input type="checkbox"/>	sodium chloride 0.9 % bolus 500 mL <i>500 mL Once As Needed IV Line Care, Starting when released, Run at TKO (25ml/hr)</i>	PRN	Route: IV
Emergency Medications		Interval	
<input type="checkbox"/>	diphenhydrAMINE (BENADRYL) injection 25-50 mg <i>25 to 50 mg Once As Needed IV Other, For mild to moderate drug reactions (flushing, dizziness, headache, diaphoresis, fever, palpitations, chest discomfort, blood pressure changes (>= 20 points in SBP), nausea, urticaria, chills, pruritis), For 1 dose, Administer 50 mg IV if patient has NOT had diphenhydramine within 2 hours of reaction. Administer 25 mg IV if patient has had diphenhydramine within 2 hours of reaction, if reaction doesn't resolve in 3 minutes may repeat 25mg IV dose for a total of 50 mg, and notify provider</i>	PRN	Route: IV

Provider Signature

EHR User ID

Date

Time

Initials

Place Patient Label Here

